



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau


INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 471/04, A61K 31/47 // (C07D 471/04, 221:00, 221:00)		A1	(11) International Publication Number: WO 98/21208
			(43) International Publication Date: 22 May 1998 (22.05.98)
(21) International Application Number: PCT/EP97/06096			
(22) International Filing Date: 5 November 1997 (05.11.97)			
(30) Priority Data: 196 46 298.3 11 November 1996 (11.11.96) DE 96118188.0 13 November 1996 (13.11.96) EP (34) Countries for which the regional or international application was filed: AT et al. 197 39 056.0 5 September 1997 (05.09.97) DE		(72) Inventor; and (75) Inventor/Applicant (for US only): FLOCKERZI, Dieter [DE/DE]; Ackerweg 26, D-78476 Allensbach (DE).	
(71) Applicant (for all designated States except US): BYK GULDEN LOMBERG CHEMISCHE FABRIK GMBH [DE/DE]; Byk-Gulden-Strasse 2, D-78467 Konstanz (DE).		(74) Common Representative: BYK GULDEN LOMBERG CHEMISCHE FABRIK GMBH; Byk-Gulden-Strasse 2, D-78467 Konstanz (DE).	
(72) Inventors (for all designated States except CA US): GUTTERER, Beate; Allensbacher Strasse 6b, D-78476 Allensbach (DE). AMSCHLER, Hermann; Hohenhewenstrasse 19, D-78315 Radolfzell (DE). ULRICH, Wolf-Rüdiger; Hebelstrasse 3, D-78464 Konstanz (DE). MARTIN, Thomas; Sonnenbühlstrasse 73, D-78464 Konstanz (DE). BÄR, Thomas; Blarerstrasse 16, D-78462 Konstanz (DE). HATZELMANN, Armin; Alter Wall 3, D-78467 Konstanz (DE). SANDERS, Karl; Mainaustrasse 207d, D-78464 Konstanz (DE). BEUME, Rolf; Bohlstrasse 13, D-78465 Konstanz (DE). BOSS, Hildegard; Flurweg 3a, D-78464		(81) Designated States: AL, AU, BA, BG, BR, CA, CN, CZ, EE, GE, HU, ID, IL, JP, KR, LT, LV, MK, MX, NO, NZ, PL, RO, SG, SI, SK, TR, UA, US, VN, YU, ZW, Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	

Published

With international search report.

(54) Title: BENZONAPHTHYRIDINES AS BRONCHIAL THERAPEUTICS



(I)

(57) Abstract

Compounds of formula (I) in which R1, R2, R3 and R4 have the meanings indicated in the description, are novel active bronchial therapeutics.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon		Republic of Korea	PT	Portugal		
CN	China	KR	Republic of Korea	RO	Romania		
CU	Cuba	KZ	Kazakhstan	RU	Russian Federation		
CZ	Czech Republic	LC	Saint Lucia	SD	Sudan		
DE	Germany	LI	Liechtenstein	SE	Sweden		
DK	Denmark	LK	Sri Lanka	SG	Singapore		
EE	Estonia	LR	Liberia				

BENZONAPHTHYRIDINES AS BRONCHIAL THERAPEUTICS

Field of application of the invention

The invention relates to novel 6-phenylbenzonaphthyridines which are used in the pharmaceutical industry for the production of medicaments.


Known technical background

DE-A 21 23 328 and USP 3,899,494 describe substituted benzonaphthyridines which are distinguished by marked inhibition of blood platelet aggregation. EP 247 971 and WO 91/17991 disclose 6-phenylbenzonaphthyridines for the treatment of inflammatory airway disorders.

Description of the invention

It has now been found that the following compounds of the formula I which are described in greater detail and differ from the compounds of EP 247 971 or WO91/17991, in particular, by the substitution on the 6-phenyl ring, have surprising and particularly advantageous properties.

The invention thus relates to compounds of the formula I



in which

R1 is 1-4C-alkyl,

R2 is hydroxyl, 1-4C-alkoxy, 3-7C-cycloalkoxy, 3-7C-cycloalkylmethoxy or 1-4C-alkoxy which is completely or predominantly substituted by fluorine,

R3 is hydroxyl, 1-4C-alkoxy, 3-7C-cycloalkoxy, 3-7C-cycloalkylmethoxy or 1-4C-alkoxy which is completely or predominantly substituted by fluorine,

or in which

R2 and R3 together are a 1-2C-alkylenedioxy group,

R4 is a phenyl radical which is substituted by R5 and R6, where

- 2 -

- R5 is hydrogen, hydroxyl, halogen, nitro, 1-4C-alkyl, trifluoromethyl or 1-4C-alkoxy,
R6 is CO-R7 or CO-R8, where
R7 is hydroxyl, 1-8C-alkoxy, 3-7C-cycloalkoxy or 3-7C-cycloalkylmethoxy and
R8 is N(R81)R82, where R81 and R82 independently of one another are hydrogen, 1-7C-alkyl, 3-7C-cycloalkyl or 3-7C-cycloalkylmethyl, or where R81 and R82, together and including the nitrogen atom to which both are bonded, are a 1-pyrrolidinyl, 1-piperidyl, 1-hexahydroazepinyl or 4-morpholinyl radical,
and to the salts of these compounds.

1-4C-Alkyl represents a straight-chain or branched alkyl radical having 1 to 4 carbon atoms. Examples which may be mentioned are the butyl, isobutyl, sec-butyl, tert-butyl, propyl, isopropyl and, preferably, the ethyl and methyl radicals.

1-4C-Alkoxy represents radicals which, in addition to the oxygen atom, contain a straight-chain or branched alkyl radical having 1 to 4 carbon atoms. Examples which may be mentioned are the butoxy, isobutoxy, sec-butoxy, tert-butoxy, propoxy, isopropoxy and, preferably, the ethoxy and methoxy radicals.

3-7C-Cycloalkoxy represents, for example, cyclopropyloxy, cyclobutyloxy, cyclopentyloxy, cyclohexyloxy and cycloheptyloxy, of which cyclopropyloxy, cyclobutyloxy and cyclopentyloxy are preferred.

3-7C-Cycloalkylmethoxy represents, for example, cyclopropylmethoxy, cyclobutylmethoxy, cyclopentylmethoxy, cyclohexylmethoxy and cycloheptylmethoxy, of which cyclopropylmethoxy, cyclobutylmethoxy and cyclopentylmethoxy are preferred.

As 1-4C-Alkoxy which is completely or predominantly substituted by fluorine, the 2,2,3,3,3-pentafluoropropoxy, the perfluoroethoxy, the 1,1,2,2-tetrafluoroethoxy, the 1,2,2-trifluoroethoxy, the trifluoromethoxy, in particular the 2,2,2-trifluoroethoxy, and preferably the difluoromethoxy radicals, for example, may be mentioned.

1-2C-Alkylenedioxy represents, for example, the methylenedioxy (-O-CH₂-O-) or the ethylenedioxy radical (-O-CH₂-CH₂-O-).

Halogen within the meaning of the invention is fluorine, chlorine or bromine.

1-8C-Alkoxy represents radicals, which, in addition to the oxygen atom, contain a straight-chain or branched alkyl radical having 1 to 8 carbon atoms. Examples which may be mentioned are the octyloxy, heptyloxy, hexyloxy, pentyloxy, methylbutoxy, ethylpropoxy, butoxy, isobutoxy, sec-butoxy, tert-

butoxy, propoxy or, preferably, the isopropoxy, ethoxy or methoxy radical.

1-7C-Alkyl represents straight-chain or branched alkyl radicals having 1 to 7 carbon atoms. Examples which may be mentioned are the heptyl, isoheptyl (5-methylhexyl), hexyl, isohexyl (4-methylpentyl), neohexyl (3,3-dimethylbutyl), pentyl, isopentyl (3-methylbutyl), neopentyl (2,2-dimethylpropyl), butyl, isobutyl, sec-butyl, tert-butyl, propyl, isopropyl, ethyl or methyl radical.

3-7C-Cycloalkyl represents the cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or cycloheptyl radical.

3-7C-Cycloalkylmethyl represents a methyl radical which is substituted by one of the abovementioned 3-7C-cycloalkyl radicals. Examples which may be mentioned are the cycloalkylmethyl radicals cyclopropylmethyl, cyclobutylmethyl and cyclopentylmethyl.

Suitable salts of compounds of the formula I - depending on substitution - are all acid addition salts or all salts with bases. The pharmacologically tolerable salts of the inorganic and organic acids and bases customarily used in pharmacy may be particularly mentioned. Those suitable are, on the one hand, water-soluble and water-insoluble acid addition salts with acids such as, for example, hydrochloric acid, hydrobromic acid, phosphoric acid, nitric acid, sulfuric acid, acetic acid, citric acid, D-gluconic acid, benzoic acid, 2-(4-hydroxybenzoyl)benzoic acid, butyric acid, sulfosalicylic acid, maleic acid, lauric acid, malic acid, fumaric acid, succinic acid, oxalic acid, tartaric acid, embonic acid, stearic acid, toluenesulfonic acid, methanesulfonic acid or 3-hydroxy-2-naphthoic acid, where the acids are employed in salt preparation - depending on whether a mono- or polybasic acid is concerned and depending on which salt is desired - in an equimolar quantitative ratio or one differing therefrom.

On the other hand - for example in the case of carboxyl substitution - salts with bases are also suitable. Examples of salts with bases which may be mentioned are alkali metal (lithium, sodium, potassium) or calcium, aluminum, magnesium, titanium, ammonium, meglumine or guanidinium salts, where here too the bases are employed in salt preparation in an equimolar quantitative ratio or one differing therefrom.

Pharmacologically intolerable salts which can be obtained first, for example, as process products in the preparation of the compounds according to the invention on an industrial scale, are converted into pharmacologically tolerable salts by methods known to the person skilled in the art.

According to expert's knowledge the compounds of the invention as well as their salts may contain, e.g. when isolated in crystalline form, varying amounts of solvents. Included within the scope of the invention are therefore all solvates and in particular all hydrates of the compounds of formula I as well as all solvates and in particular all hydrates of the salts of the compounds of formula I.

Compounds of the formula I to be emphasized are those in which

- R1 is 1-4C-alkyl,
- R2 is 1-4C-alkoxy, 3-7C-cycloalkoxy, 3-7C-cycloalkylmethoxy or 1-2C-alkoxy which is completely or predominantly substituted by fluorine,
- R3 is 1-4C-alkoxy, 3-7C-cycloalkoxy, 3-7C-cycloalkylmethoxy or 1-2C-alkoxy which is completely or predominantly substituted by fluorine,
- R4 is a phenyl radical which is substituted by R5 and R6, where
- R5 is hydrogen, hydroxyl, halogen, nitro, 1-4C-alkyl, trifluoromethyl or 1-4C-alkoxy,
- R6 is CO-R7 or CO-R8, where
- R7 is hydroxyl, 1-8C-alkoxy, 3-7C-cycloalkoxy or 3-7C-cycloalkylmethoxy and
- R8 is N(R81)R82, where R81 and R82 independently of one another are hydrogen, 1-7C-alkyl, 3-7C-cycloalkyl or 3-7C-cycloalkylmethyl, or where R81 and R82, together and including the nitrogen atom to which both are bonded, are a 1-piperidyl, 1-hexahydroazepinyl or 4-morpholinyl radical,

and the salts of these compounds.

One embodiment of the compounds of the formula I to be emphasized are those compounds in which

- R1 is 1-4C-alkyl,
- R2 is 1-4C-alkoxy, 3-7C-cycloalkoxy, 3-7C-cycloalkylmethoxy or 1-2C-alkoxy which is completely or predominantly substituted by fluorine,
- R3 is 1-4C-alkoxy, 3-7C-cycloalkoxy, 3-7C-cycloalkylmethoxy or 1-2C-alkoxy which is completely or predominantly substituted by fluorine,
- R4 is a phenyl radical which is substituted by R5 and R6, where
- R5 is hydrogen, hydroxyl, halogen, nitro, 1-4C-alkyl, trifluoromethyl or 1-4C-alkoxy,
- R6 is CO-R7 or CO-R8, where
- R7 is hydroxyl, 1-8C-alkoxy, 3-7C-cycloalkoxy or 3-7C-cycloalkylmethoxy and
- R8 is N(R81)R82, where R81 and R82 independently of one another are hydrogen, 1-7C-alkyl, 3-7C-cycloalkyl or 3-7C-cycloalkylmethyl, or where R81 and R82, together and including the nitrogen atom to which both are bonded, are a 1-piperidyl or 1-hexahydroazepinyl radical,

and the salts of these compounds.

Compounds of the formula I particularly to be emphasized are those in which

- R1 is methyl,
- R2 is 1-4C-alkoxy, 3-7C-cycloalkoxy or 1-2C-alkoxy which is completely or predominantly substituted by fluorine,
- R3 is 1-4C-alkoxy, 3-7C-cycloalkoxy or 1-2C-alkoxy which is completely or predominantly substituted by fluorine,

- 5 -

R4 is a phenyl radical which is substituted by R5 and R6, where
R5 is hydrogen, hydroxyl, 1-4C-alkyl or 1-4C-alkoxy,
R6 is CO-R7 or CO-R8, where
R7 is hydroxyl, 1-8C-alkoxy or 3-7C-cycloalkoxy and
R8 is N(R81)R82, where R81 and R82 independently of one another are hydrogen, 1-7C-alkyl or 3-7C-cycloalkyl, or where R81 and R82, together and including the nitrogen atom to which both are bonded, are a 1-piperidyl, 1-hexahydroazepinyl or 4-morpholinyl radical,
and the salts of these compounds.

One embodiment of the compounds of the formula I particularly to be emphasized are those compounds in which

R1 is methyl,
R2 is 1-4C-alkoxy, 3-7C-cycloalkoxy or 1-2C-alkoxy which is completely or predominantly substituted by fluorine,
R3 is 1-4C-alkoxy, 3-7C-cycloalkoxy or 1-2C-alkoxy which is completely or predominantly substituted by fluorine,
R4 is a phenyl radical which is substituted by R5 and R6, where
R5 is hydrogen, hydroxyl, 1-4C-alkyl or 1-4C-alkoxy,
R6 is CO-R7 or CO-R8, where
R7 is hydroxyl, 1-8C-alkoxy or 3-7C-cycloalkoxy and
R8 is N(R81)R82, where R81 and R82 independently of one another are hydrogen, 1-7C-alkyl or 3-7C-cycloalkyl,
and the salts of these compounds.

Preferred compounds of the formula I are those in which

R1 is methyl,
R2 is methoxy or ethoxy,
R3 is methoxy or ethoxy,
R4 is a phenyl radical which is substituted by R5 and R6, where
R5 is hydrogen,
R6 is CO-R7 or CO-R8, where
R7 is hydroxyl or 1-8C-alkoxy and
R8 is N(R81)R82, where R81 and R82 independently of one another are hydrogen or 1-4C-alkyl or 5-7C-cycloalkyl, or where R81 and R82, together and including the nitrogen atom to which both are bonded, are a 1-piperidyl, 1-hexahydroazepinyl or 4-morpholinyl radical,
and the salts of these compounds.

One embodiment of the preferred compounds of the formula I are those compounds in which

- 6 -


- R1 is methyl,
 R2 is methoxy or ethoxy,
 R3 is methoxy or ethoxy,
 R4 is a phenyl radical which is substituted by R5 and R6, where
 R5 is hydrogen,
 R6 is CO-R7 or CO-R8, where
 R7 is hydroxyl or 1-8C-alkoxy and
 R8 is N(R81)R82, where R81 and R82 independently of one another are hydrogen or 1-4C-alkyl,
 and the salts of these compounds.

Particularly preferred compounds of the formula I are those in which

- R1 is methyl,
 R2 is ethoxy,
 R3 is methoxy or ethoxy,
 R4 is a phenyl radical which is substituted by R5 and R6, where
 R5 is hydrogen,
 R6 is CO-R7 or CO-R8, where
 R7 is 1-4C-alkoxy and
 R8 is N(R81)R82, where R81 and R82 independently of one another are 1-4C-alkyl or 5-7C-cycloalkyl, or where R81 and R82, together and including the nitrogen atom to which both are bonded, are a 1-piperidyl or 1-hexahydroazepinyl radical,
 and the salts of these compounds.

The compounds of the formula I are chiral compounds having chiral centers in positions 4a and 10b

Numbering:



The invention therefore includes all conceivable pure diastereomers and pure enantiomers and mixtures thereof in any mixing ratio, including the racemates. Preferred are those compounds of the formula I in which the hydrogen atoms in positions 4a and 10b are in the cis position relative to one another. The pure cis enantiomers and their mixtures in any mixing ratio and including the racemates are particularly preferred here.

Particularly preferred in this connection are those compounds of the formula I which in positions 4a and 10b have the same absolute configuration as the compound (-)-cis-4-amino-3-(3,4-dimethoxyphenyl)-1-methylpiperidine dihydrochloride having the optical rotation $[\alpha]_D^{22} = -57.1^\circ$ (c=1, methanol) which can be employed as a starting material and is described in DE 42 17 401.

Especially to be emphasized are those compounds of formula I which can be prepared starting from the starting compounds (-)-cis-4-Amino-3-(3,4-diethoxyphenyl)-1-methylpiperidine dihydrochloride (example F) or (-)-cis-4-Amino-3-(3-ethoxy-4-methoxyphenyl)-1-methylpiperidine dihydrochloride (example O) and are selected from

cis-8,9-Diethoxy-6-(4-isopropoxycarbonylphenyl)-2-methyl-1,2,3,4,4a,10b-hexahydrobenzo[c][1,6]-naphthyridine;

cis-8,9-Diethoxy-6-(4-diisopropylaminocarbonylphenyl)-2-methyl-1,2,3,4,4a,10b-hexahydrobenzo[c][1,6]naphthyridine;

cis-8,9-Diethoxy-6-(4-N-cyclohexyl-N-isopropylaminocarbonylphenyl)-2-methyl-1,2,3,4,4a,10b-hexahydrobenzo[c][1,6]naphthyridine;

cis-8,9-Diethoxy-6-(4-dibutylaminocarbonylphenyl)-2-methyl-1,2,3,4,4a,10b-hexahydrobenzo[c][1,6]naphthyridine;

cis-8,9-Diethoxy-6-[4-(hexahydroazepin-1-yl-carbonyl)phenyl]-2-methyl-1,2,3,4,4a,10b-hexahydrobenzo[c][1,6]naphthyridine;

cis-8,9-Diethoxy-6-[4-(piperidin-1-ylcarbonyl)phenyl]-2-methyl-1,2,3,4,4a,10b-hexahydrobenzo[c][1,6]naphthyridine;

cis-9-Ethoxy-8-methoxy-6-(4-diisopropylaminocarbonylphenyl)-2-methyl-1,2,3,4,4a,10b-hexahydrobenzo[c][1,6]naphthyridine;

cis-9-Ethoxy-8-methoxy-6-(4-dibutylaminocarbonylphenyl)-2-methyl-1,2,3,4,4a,10b-hexahydrobenzo[c][1,6]naphthyridine;

cis-9-Ethoxy-8-methoxy-6-[4-(hexahydroazepin-1-yl-carbonyl)phenyl]-2-methyl-1,2,3,4,4a,10b-hexahydrobenzo[c][1,6]naphthyridine;


cis-9-Ethoxy-8-methoxy-6-[4-(piperidin-1-yl-carbonyl)phenyl]-2-methyl-1,2,3,4,4a,10b-hexahydrobenzo[c][1,6]naphthyridine

and the salts of these compounds.

The enantiomers can be separated in a known manner (for example by preparation and separation of corresponding diastereoisomeric compounds) or by stereoselective synthesis methods. Such separation processes and synthesis methods are described, for example, in EP 247 971 and in DE 42 17 401.

The invention further relates to a process for the preparation of the compounds of the formula I, in which R1, R2, R3 and R4 have the meanings indicated above, and their salts. The process comprises

subjecting compounds of the formula II




in which R1, R2, R3 and R4 have the meanings indicated above, to a cyclocondensation reaction and, if desired, then converting the compounds of the formula I obtained into their salts, or, if desired, then converting salts of the compounds of the formula I obtained into the free compounds.

If desired, compounds of the formula I obtained can be converted into further compounds of the formula I by derivatization. For example, the corresponding acids can be obtained from compounds of the formula I in which R4 is a phenyl radical which is substituted by R5 and R6, and R6 is an ester group, by acidic or alkaline hydrolysis, or the corresponding amides can be prepared by reaction with amines of the formula HN(R81)R82, in which R81 and R82 have the meanings indicated above. The reactions are expediently carried out analogously to the methods known to the person skilled in the art, e.g. as described in the following examples.

The cyclocondensation is carried out in a manner known per se to the person skilled in the art according to Bischler-Napieralski (e.g. as described in J. Chem. Soc., 1956, 4280-4282) in the presence of a suitable condensing agent, such as, for example, polyphosphoric acid, phosphorus pentachloride, phosphorus trichloride, phosphorus pentoxide, thionyl chloride or preferably phosphorus oxytrichloride, in a suitable inert solvent, e.g. in a chlorinated hydrocarbon such as chloroform, or in a cyclic hydrocarbon such as toluene or xylene, or another inert solvent such as acetonitrile, or without a further solvent using an excess of condensing agent, preferably at elevated temperature, in particular at the boiling temperature of the solvent or condensing agent used.

Compounds of the formula II in which R1, R2, R3 and R4 have the meanings indicated above are accessible from the corresponding compounds of the formula III

- 9 -



in which R1, R2 and R3 have the meanings indicated above, by reaction with compounds of the formula R4-CO-X in which R4 has the meaning indicated above and X is a suitable leaving group, preferably a chlorine atom. For example, the benzylation is carried out as in the following examples according to the Einhorn process, the Schotten-Baumann variant or as described in J. Chem. Soc. (C), 1971, 1805-1808.

The preparation of cis/trans racemate mixtures and of pure cis racemates of compounds of the formula III is described, for example, in US patent 3,899,494, in DE-A 21 23 328 and in DE-A 16 95 782. Pure cis enantiomers of the compounds of the formula III can be obtained, for example, by the processes as are disclosed in EP 0 247 971 and in DE 42 17 401.

Compounds of the formula R4-CO-X are either known or can be prepared in a known manner.

The substances according to the invention are isolated and purified in a manner known per se, for example by distilling off the solvent in vacuo and recrystallizing the residue obtained from a suitable solvent or subjecting it to one of the customary purification methods, such as, for example, column chromatography on a suitable support material.

Salts are obtained by dissolving the free compound in a suitable solvent, e.g. in a chlorinated hydrocarbon, such as methylene chloride or chloroform, or a low molecular weight aliphatic alcohol (ethanol, isopropanol) which contains the desired acid or base, or to which the desired acid or base is then added. The salts are obtained by filtering, reprecipitating, precipitating with a nonsolvent for the addition salt or by evaporating the solvent. Salts obtained can be converted into the free compounds, which can in turn be converted into salts, by alkalization or by acidification. In this manner, pharmacologically intolerable salts can be converted into pharmacologically tolerable salts.

The following examples serve to illustrate the invention in greater detail without restricting it. Further compounds of the formula I, whose preparation is not explicitly described, can also be prepared in an analogous manner or in a manner familiar per se to the person skilled in the art using customary process techniques.

- 10 -

In the examples, m.p. stands for melting point, h for hour(s), RT for room temperature, EF for empirical formula, MW for molecular weight, calc. for calculated. The compounds mentioned in the examples and their salts are a preferred subject of the invention.

ExamplesFinal products

1. (-)-cis-8,9-Dimethoxy-6-(4-methoxycarbonylphenyl)-2-methyl-1,2,3,4,4a,10b-hexahydro-benzo[c][1,6]naphthyridine hydrochloride

2.31 g of (-)-cis-terephthalic acid N-[3-(3,4-dimethoxyphenyl)-1-methylpiperidin-4-yl]amide mono-methyl ester are heated to boiling under reflux for 4 h in 25 ml of acetonitrile and 3 ml of phosphorus oxytrichloride. After distilling off the excess phosphorus oxytrichloride, the residue is partitioned between dichloromethane and saturated sodium hydrogencarbonate solution. The organic phase is washed with water, dried over sodium sulfate and concentrated. The solid residue is purified by silica gel chromatography, the main product fraction is separated and this is concentrated. The solid residue is dissolved in a little methanol, and the solution is treated with one equivalent of aqueous HCl and concentrated. The solid residue is recrystallized in methanol/diethyl ether. 1.76 g (70% of theory) of the title compound are obtained as a 1.25-hydrochloride 0.5-hydrate having the m.p. 188-192°C (unsharp, slow deliquescence).

EF: $C_{23}H_{26}N_2O_4 \times 1.25 HCl \times 0.5 H_2O$; MW: 449.05

Elemental analysis: calc.: C 61.52 H 6.34 Cl 9.87 N 6.24

found: C 61.52 H 6.19 Cl 9.93 N 6.23

Optical rotation: $[\alpha]_D^{20} = -66^\circ$ (c=1, methanol).

Starting from the corresponding starting compounds described below, the following title compounds are obtained analogously to Example 1:

2. (-)-cis-8,9-Dimethoxy-6-(3-methoxycarbonylphenyl)-2-methyl-1,2,3,4,4a,10b-hexahydro-benzo[c][1,6]naphthyridine hydrochloride

EF: $C_{23}H_{26}N_2O_4 \times 1.1 HCl \times 0.23 H_2O$; MW: 438.69; m.p. from about 155°C slow deliquescence; yield: 63% of theory

Elemental analysis: calc.: C 63.08 H 6.32 Cl 8.90 N 6.40

found: C 63.13 H 6.53 Cl 8.81 N 6.53

Optical rotation: $[\alpha]_D^{20} = -90.8^\circ$ (c=1, methanol).

3. (-)-cis-8,9-Dimethoxy-6-(4-isopropoxycarbonylphenyl)-2-methyl-1,2,3,4,4a,10b-hexahydrobenzo[c][1,6]naphthyridine hydrochloride

EF: $C_{25}H_{30}N_2O_4 \times 1.15 \text{ HCl} \times 0.8 \text{ H}_2\text{O}$; MW: 478.79; m.p. 166-170°C; yield: 65% of theory;

Elemental analysis: calc.: C 62.84 H 6.88 Cl 8.53 N 5.86

found: C 62.92 H 7.06 Cl 8.44 N 6.04

Optical rotation: $[\alpha]_D^{20} = -40.4^\circ$ (c=1, methanol).

4. (-)-cis-8,9-Diethoxy-6-(4-methoxycarbonylphenyl)-2-methyl-1,2,3,4,4a,10b-hexahydrobenzo[c][1,6]naphthyridine hydrochloride

EF: $C_{25}H_{30}N_2O_4 \times 1.25 \text{ HCl} \times 1.12 \text{ H}_2\text{O}$; MW: 485.9; m.p. 143-148°C; yield: 76% of theory;

Elemental analysis: calc.: C 61.79 H 6.97 Cl 8.18 N 5.76

found: C 61.88 H 6.88 Cl 8.34 N 5.70

Optical rotation: $[\alpha]_D^{20} = -50.2^\circ$ (c=1, methanol).

5. (-)-cis-8,9-Diethoxy-6-(4-isopropoxycarbonylphenyl)-2-methyl-1,2,3,4,4a,10b-hexahydrobenzo[c][1,6]naphthyridine hydrochloride

EF: $C_{27}H_{34}N_2O_4 \times 1.1 \text{ HCl} \times 0.83 \text{ H}_2\text{O}$; MW: 505.64; m.p. 205-209°C, from about 165°C slow agglutination; yield: 69% of theory;

Elemental analysis: calc.: C 64.17 H 7.33 Cl 7.72 N 5.65

found: C 64.29 H 7.37 Cl 7.60 N 5.65

Optical rotation: $[\alpha]_D^{20} = -47.1^\circ$ (c=1, methanol).

6. (-)-cis-6-(4-Aminocarbonylphenyl)-8,9-dimethoxyphenyl-2-methyl-1,2,3,4,4a,10b-hexahydrobenzo[c][1,6]naphthyridine

The title compound from Example 1 is allowed to stand at RT for 48 h in a 1+1 vol. mixture of methanol and conc. ammonia solution. After concentrating completely, the solid residue is recrystallized in a mixture of 1 part by volume of methanol and 10 parts by volume of diethyl ether. 51% of theory of the title compound of m.p. 229-232°C is obtained.

EF: $C_{22}H_{25}N_3O_3$; MW: 379.46;

Elemental analysis: calc.: C 69.64 H 6.64 N 11.07

found: C 69.41 H 6.54 N 11.00

Optical rotation: $[\alpha]_D^{20} = -104.7^\circ$ (c=1, methanol).

7. (-)-cis-6-(4-Carboxyphenyl)-8,9-dimethoxyphenyl-2-methyl-1,2,3,4,4a,10b-hexahydro-benzo[c][1,6]naphthyridine

The ester group of the title compound from Example 1 is hydrolyzed at RT in a mixture of methanol and 2N sodium hydroxide solution. After hydrolysis is complete, the mixture is neutralized by addition of an appropriate amount of hydrochloric acid and the suspension obtained is largely concentrated (removal of the methanol). The solid residue is suspended using water and filtered off with suction. After thorough washing with water, the filter residue consists of the title compound, which after drying has an m.p. of 237-240°C.

EF: $C_{22}H_{24}N_2O_4 \times 0.5 H_2O$; MW: 389.46; yield: 80% of theory;

Elemental analysis: calc.: C 67.85 H 6.47 N 7.19

found: C 68.05 H 6.62 N 7.24

Analogously to Example 7, the following 2 title compounds are obtained by hydrolysis of the corresponding esters described above:

8. (-)-cis-6-(3-Carboxyphenyl)-8,9-dimethoxyphenyl-2-methyl-1,2,3,4,4a,10b-hexahydro-benzo[c][1,6]naphthyridine

EF: $C_{22}H_{24}N_2O_4$; MW: 380.45; yield: 76% of theory; m.p. from about 168° slow agglutination, from about 230°C decomposition.

9. (-)-cis-6-(4-Carboxyphenyl)-8,9-diethoxyphenyl-2-methyl-1,2,3,4,4a,10b-hexahydro-benzo[c][1,6]naphthyridine

EF: $C_{24}H_{28}N_2O_4$; MW: 408.50; yield: 88% of theory; m.p. >240°C (decomposition).

The following title compounds are obtained analogously to Example 1 when the piperidines appropriately substituted in the 4-position are employed as starting compounds for the cyclocondensation. These are obtained in the manner described below for the starting compound A when appropriate terephthalic acid monoamides are employed.

10. (-)-cis-8,9-Diethoxy-6-(4-dimethylaminocarbonylphenyl)-2-methyl-1,2,3,4,4a,10b-hexahydrobenzo[c][1,6]naphthyridine hydrochloride

EF: $C_{26}H_{33}N_3O_3 \times HCl \times H_2O$; MW: 490.05; m.p.: 142-150°C (solid foamed product); yield: 46% of theory;

Elemental analysis: calc.: C 63.73 H 7.40 Cl 7.23 N 8.57

found: C 64.08 H 7.32 Cl 7.48 N 8.31

Optical rotation: $[\alpha]_D^{20} = -28.9^\circ$ (c=1, methanol).

11. (-)-cis-8,9-Diethoxy-6-(4-diisopropylaminocarbonylphenyl)-2-methyl-1,2,3,4,4a,10b-hexahydrobenzo[c][1,6]naphthyridine hydrochloride

EF: $C_{30}H_{41}N_3O_3 \times HCl \times 1.4 H_2O$; MW: 553.37; m.p.: 164-180°C (unsharp range); yield: 30% of theory;

Elemental analysis: calc.: C 65.12 H 8.16 Cl 6.41 N 7.59

found: C 64.85 H 8.29 Cl 6.50 N 7.66

Optical rotation: $[\alpha]_D^{20} = -42.0^\circ$ (c=1, methanol).

12. (-)-cis-8,9-Dimethoxy-6-(4-diisopropylaminocarbonylphenyl)-2-methyl-1,2,3,4,4a,10b-hexahydrobenzo[c][1,6]naphthyridine hydrochloride

EF: $C_{28}H_{37}N_3O_3 \times HCl \times 2.4 H_2O$; MW: 542.32; m.p.: 175-185°C (unsharp range); yield: 24% of theory;

Elemental analysis: calc.: C 62.01 H 7.77 Cl 6.54 N 7.75

found: C 61.88 H 7.81 Cl 6.68 N 7.73

Optical rotation: $[\alpha]_D^{20} = -60.6^\circ$ (c=1, methanol).

13. (-)-cis-8,9-Dimethoxy-6-(4-dimethylaminocarbonylphenyl)-2-methyl-1,2,3,4,4a,10b-hexahydrobenzo[c][1,6]naphthyridine hydrochloride

EF: $C_{24}H_{29}N_3O_3 \times HCl \times 0.6 H_2O$; MW: 454.79; m.p.: 226-228°C (decomposition); yield: 21% of theory;

Elemental analysis: calc.: C 63.38 H 6.92 Cl 7.80 N 9.24

found: C 63.10 H 7.13 Cl 8.12 N 9.14

Optical rotation: $[\alpha]_D^{20} = -58.0^\circ$ (c=1, methanol).

14. (+)-cis-8,9-Diethoxy-2-methyl-6-[4-(4-morpholinocarbonyl)phenyl]-1,2,3,4,4a,10b-hexahydrobenzo[c][1,6]naphthyridine hydrochloride

EF: $C_{28}H_{35}N_3O_4 \times 1.25 HCl \times H_2O$; MW: 541.20; m.p.: 165-170°C (unsharp range); yield: 45% of the-ory;

Elemental analysis: calc.: C 62.14 H 7.12 Cl 8.19 N 7.76

found: C 62.30 H 7.21 Cl 7.96 N 7.35

Optical rotation: $[\alpha]_D^{20} = +12.4^\circ$ (c=1, methanol).

15. (+)-cis-8,9-Diethoxy-6-(4-dicyclohexylaminocarbonylphenyl)-2-methyl-1,2,3,4,4a,10b-hexahydrobenzo[c][1,6]naphthyridine hydrochloride

EF: $C_{36}H_{49}N_3O_3 \times 1.2 HCl \times 1.4 H_2O$; MW: 640.78; m.p.: 195-202°C (unsharp range); yield: 65% of theory;

Elemental analysis: calc.: C 67.48 H 8.34 Cl 6.64 N 6.56

found: C 67.53 H 8.16 Cl 6.62 N 6.71

Optical rotation: $[\alpha]_D^{20} = +24.2^\circ$ (c=1, methanol).

16. (+)-cis-8,9-Diethoxy-6-(4-N-cyclohexyl-N-isopropylaminocarbonylphenyl)-2-methyl-1,2,3,4,4a,10b-hexahydrobenzo[c][1,6]naphthyridine hydrochloride

EF: $C_{33}H_{45}N_3O_3 \times 1.25 HCl \times 1 H_2O$; MW: 595.33; m.p.: 163-195°C (unsharp range); yield: 57% of theory;

Elemental analysis: calc.: C 66.58 H 8.17 Cl 7.44 N 7.06

found: C 66.71 H 8.04 Cl 7.42 N 7.25

Optical rotation: $[\alpha]_D^{20} = +21.6^\circ$ (c=1, methanol).

17. (+)-cis-8,9-Diethoxy-6-(4-dibutylaminocarbonylphenyl)-2-methyl-1,2,3,4,4a,10b-hexahydrobenzo[c][1,6]naphthyridine hydrochloride

EF: $C_{32}H_{45}N_3O_3 \times 2 HCl \times 0.9 H_2O$; MW: 608.88; m.p.: 144-162°C (unsharp range; decomposition); yield: 62% of theory;

Elemental analysis: calc.: C 63.13 H 8.08 Cl 11.65 N 6.90

found: C 63.29 H 8.18 Cl 11.61 N 6.80

Optical rotation: $[\alpha]_D^{20} = +191.7^\circ$ (c=1, methanol).

18. (-)-cis-8,9-Diethoxy-6-[4-(hexahydroazepin-1-yl-carbonyl)phenyl]-2-methyl-1,2,3,4,4a,10b-hexahydrobenzo[c][1,6]naphthyridine hydrochloride

EF: $C_{30}H_{39}N_3O_3 \times 1.2 \text{ HCl} \times 0.6 \text{ H}_2\text{O}$; MW: 544.23; m.p.: 138-154°C (unsharp range); yield: 58% of theory;

Elemental analysis: calc.: C 66.21 H 7.66 Cl 7.81 N 7.72

found: C 66.17 H 7.70 Cl 7.80 N 7.71

Optical rotation: $[\alpha]_D^{20} = -11.5^\circ$ (c=1, methanol).

19. (-)-cis-9-Ethoxy-8-methoxy-6-(4-diisopropylaminocarbonylphenyl)-2-methyl-1,2,3,4,4a,10b-hexahydrobenzo[c][1,6]naphthyridine hydrochloride

EF: $C_{29}H_{39}N_3O_3 \times \text{HCl} \times 0.6 \text{ H}_2\text{O}$; MW: 524.92; m.p.: 175-179°C (unsharp); yield: 67% of theory;

Elemental analysis: calc.: C 66.36 H 7.91 Cl 6.75 N 8.01

found: C 66.28 H 7.99 Cl 6.87 N 7.97

Optical rotation: $[\alpha]_D^{20} = -42.7^\circ$ (c=1, methanol).

20. (-)-cis-8-Ethoxy-9-methoxy-6-(4-diisopropylaminocarbonylphenyl)-2-methyl-1,2,3,4,4a,10b-hexahydrobenzo[c][1,6]naphthyridine hydrochloride

EF: $C_{29}H_{39}N_3O_3 \times 1.03 \text{ HCl} \times 0.94 \text{ H}_2\text{O}$; MW: 532.16; m.p.: 176-179°C (unsharp); yield: 43% of theory;

Elemental analysis: calc.: C 65.49 H 7.94 Cl 6.87 N 7.90

found: C 65.43 H 7.71 Cl 6.86 N 7.99

Optical rotation: $[\alpha]_D^{20} = -48.0^\circ$ (c=1, methanol).

21. (-)-cis-9-Ethoxy-8-methoxy-6-(4-methoxycarbonylphenyl)-2-methyl-1,2,3,4,4a,10b-hexahydrobenzo[c][1,6]naphthyridine hydrochloride

EF: $C_{24}H_{28}N_2O_4 \times 1.05 \text{ HCl} \times 1.27 \text{ H}_2\text{O}$; MW:; m.p.: 150-160°C (sintering, unsharp range); yield: 89% of theory;

Elemental analysis: calc.: C 61.37 H 6.78 Cl 7.92 N 5.96

found: C 61.39 H 6.77 Cl 7.93 N 5.94

Optical rotation: $[\alpha]_D^{20} = -84.3^\circ$ (c=1, methanol).

22. (-)-cis-9-Ethoxy-8-methoxy-6-(3-methoxycarbonylphenyl)-2-methyl-1,2,3,4,4a,10b-hexahydrobenzo[c][1,6]naphthyridine hydrochloride

EF: $C_{24}H_{28}N_2O_4 \times \text{HCl} \times 0.66 \text{ H}_2\text{O}$; MW: 456.82; m.p.: sintering from about 140°C, slow melting with

decomposition up to about 150°C; yield: 88% of theory;

Elemental analysis: calc.: C 63.11 H 6.69 Cl 7.76 N 6.13

found: C 62.98 H 6.78 Cl 7.89 N 6.07

Optical rotation: $[\alpha]_D^{20} = -143.1^\circ$ (c=1, methanol).

23. (-)-cis-9-Ethoxy-8-methoxy-6-(4-dibutylaminocarbonylphenyl)-2-methyl-1,2,3,4,4a,10b-hexahydrobenzo[c][1,6]naphthyridine hydrochloride

EF: $C_{31}H_{43}N_3O_3 \times 1.1 \text{ HCl} \times 1.17 \text{ H}_2\text{O}$; MW: 566.82; m.p.: 104-112°C (solid foamed product, slow deliquescence); yield: 60% of theory;

Elemental analysis: calc.: C 65.68 H 8.26 Cl 6.88 N 7.41

found: C 65.80 H 8.09 Cl 6.97 N 7.49

Optical rotation: $[\alpha]_D^{20} = -16.2^\circ$ (c=1, methanol).

24. (-)-cis-9-Ethoxy-8-methoxy-6-[4-(hexahydroazepin-1-yl-carbonyl)phenyl]-2-methyl-1,2,3,4,4a,10b-hexahydrobenzo[c][1,6]naphthyridine

EF: $C_{29}H_{37}N_3O_3$; MW: 475.65; m.p.: 136-142°C (unsharp range, sintering from about 125°C); yield: 66% of theory;

Elemental analysis: calc.: C 73.23 H 7.84 N 8.83

found: C 73.02 H 8.08 N 8.67

Optical rotation: $[\alpha]_D^{20} = -72.5^\circ$ (c=1, methanol).

25. (-)-cis-9-Ethoxy-8-methoxy-6-[4-(piperidin-1-yl-carbonyl)phenyl]-2-methyl-1,2,3,4,4a,10b-hexahydrobenzo[c][1,6]naphthyridine hydrochloride

EF: $C_{28}H_{25}N_3O_3 \times 0.4 \text{ H}_2\text{O}$; MW: 468.82; m.p.: 79-82°C (unsharp range); yield: 43% of theory;

Elemental analysis: calc.: C 71.70 H 7.70 N 8.96

found: C 71.78 H 7.71 N 8.97

Optical rotation: $[\alpha]_D^{20} = -77.2^\circ$ (c=1, methanol).

Analogously to Example 7, the following 2 title compounds are obtained by hydrolysis of the corresponding esters described above.

26. (-)-cis-6-(4-carboxyphenyl)-9-ethoxy-8-methoxyphenyl-2-methyl-1,2,3,4,4a,10b-hexahydrobenzo[c][1,6]naphthyridine

EF: $C_{23}H_{26}N_2O_4 \times 1.1 \text{ H}_2\text{O}$; MW: 414.29; m.p.: 240-242°C (decomposition with red coloration); yield:

- 18 -

91% of theory;

Elemental analysis: calc.: C 66.68 H 6.86 N 6.76

found: C 66.81 H 6.75 N 6.70

Optical rotation: $[\alpha]_D^{20} = -109.7^\circ$ (c=1, methanol + 1.0 equivalent of 0.1N NaOH).

27. (-)-cis-6-(3-Carboxyphenyl)-9-ethoxy-8-methoxyphenyl-2-methyl-1,2,3,4,4a,10b-hexahydrobenzo[c][1,6]naphthyridine

EF: $C_{23}H_{26}N_2O_4 \times 0.54 H_2O$; MW: 404.17; m.p.: 158-170°C (decomposition); yield: 83% of theory;

Elemental analysis: calc.: C 68.36 H 6.75 N 6.93

found: C 68.25 H 6.86 N 6.96

Optical rotation: $[\alpha]_D^{20} = -150.7^\circ$ (c=1, methanol + 1.0 equivalent of 0.1N NaOH).

Starting compounds

A. (-)-cis-Terephthalic acid N-[3-(3,4-dimethoxyphenyl)-1-methylpiperidin-4-yl]amide monomethyl ester

A solution of monomethyl terephthaloyl chloride (prepared from 2.2 g of monomethyl terephthalate and thionyl chloride) in 5 ml of dichloromethane is added dropwise at RT in the course of 10 min. to a solution of 3 g of (-)-cis-4-amino-3-(3,4-dimethoxyphenyl)-1-methylpiperidine (prepared by extraction of the free base with dichloromethane after treatment of the corresponding dihydrochloride ($[\alpha]_D^{20} = -57.1^\circ$, c=1, methanol) with dilute sodium hydroxide solution) in 10 ml of dichloromethane and 1.9 ml of triethylamine. After stirring for about 2 h, the mixture is extracted with about 50 ml of saturated sodium hydrogencarbonate solution, and the organic phase is washed a further two times with 50 ml of water each time and dried over sodium sulfate. The viscous residue remaining after concentration is purified by column chromatography. The main product fraction concentrated in vacuo affords a solid foaming residue which is recrystallized in a mixture of methanol and diethyl ether (about 1 + 1 vol.). 2.3 g of the title compound having the m.p. 151-152°C are obtained; yield: 47% of theory; EF: $C_{23}H_{28}N_2O_5$; MW: 412.48.

Elemental analysis: calc. C 66.97 H 6.84 N 6.79

found C 66.82 H 6.97 N 6.97

Optical rotation: $[\alpha]_D^{22} = -74.9^\circ$ (c=1, methanol).

Starting from the corresponding starting compounds, the following title compounds are obtained analogously to the procedure according to Example A:

B. (-)-cis-Isophthalic acid N-[3-(3,4-dimethoxyphenyl)-1-methylpiperidin-4-yl]amide mono-methyl ester

EF: C₂₃H₂₈N₂O₅; MW: 412.49; yield: 63% of theory; m.p.: 122-123°C;

Optical rotation: $[\alpha]_D^{20} = -30.6^\circ$ (c=1, methanol).

C. (-)-cis-Terephthalic acid N-[3-(3,4-dimethoxyphenyl)-1-methylpiperidin-4-yl]amide monoisopropyl ester

EF: C₂₅H₃₂N₂O₅; MW: 440.54; yield: 60% of theory; m.p.: 136-142°C (unsharp);

Optical rotation: $[\alpha]_D^{20} = -48.2^\circ$ (c=1, methanol).

D. (-)-cis-Terephthalic acid N-[3-(3,4-diethoxyphenyl)-1-methylpiperidin-4-yl]amide mono-methyl ester

EF: C₂₅H₃₂N₂O₅; MW: 440.54; yield: 55% of theory; m.p.: 169-173°C (unsharp);

Optical rotation: $[\alpha]_D^{20} = -66.4^\circ$ (c=1, methanol).

The title compound is obtained by the process described in Example A when (-)-cis-4-amino-3-(3,4-diethoxyphenyl)-1-methylpiperidine having the optical rotation $[\alpha]_D^{20} = -35.1^\circ$ (dihydrochloride, solid foamed crude product, c=1, methanol) is employed as the amine component.

E. (-)-cis-Terephthalic acid N-[3-(3,4-diethoxyphenyl)-1-methylpiperidin-4-yl]amide mono-isopropyl ester

EF: C₂₇H₃₆N₂O₅; MW: 468.6; yield: 63% of theory; m.p.: 119-126°C (unsharp);

Optical rotation: $[\alpha]_D^{20} = -51.5^\circ$ (c=1, methanol).

F. (-)-cis-4-Amino-3-(3,4-diethoxyphenyl)-1-methylpiperidine dihydrochloride

The title compound is obtained analogously to the process described in DE 42 17 401 when the corresponding 3,4-diethoxy compounds are employed in the examples described there.

EF: C₁₆H₂₆N₂O₂ × 2HCl; MW: 351.32; obtained as a solid foamed crude product; m.p.: shrinkage and slow deliquescence from about 120°C, unsharp melting range up to about 150°C;

Optical rotation: $[\alpha]_D^{20} = -35.1^\circ$ (c=1, methanol).

G. (-)-cis-N-[3-(3,4-Diethoxyphenyl)-1-methylpiperidin-4-yl]-4-(morpholine-4-carbonyl)benzamide

EF: $C_{28}H_{37}N_3O_5$; MW: 495.6; yield: 71% of theory; m.p.: 178-179°C;

Optical rotation: $[\alpha]_D^{20} = -57.3^\circ$ (c=1, methanol).

H. (-)-cis-Terephthalic acid N,N-dibutyl-N'-[3-(3,4-diethoxyphenyl)-1-methylpiperidin-4-yl]diamide

EF: $C_{32}H_{47}N_3O_4 \times 0.25 H_2O$; MW: 537.75; yield: 76% of theory; m.p.: 115-120°C;

Optical rotation: $[\alpha]_D^{20} = -57.0^\circ$ (c=1, methanol).

I. (-)-cis-Terephthalic acid N-cyclohexyl-N'-[3-(3,4-diethoxyphenyl)-1-methylpiperidin-4-yl]-N-isopropylidiamide

EF: $C_{33}H_{47}N_3O_4$; MW: 549.76; yield: 66% of theory; m.p.: 59-64°C (unsharp range, solidified foam);

Optical rotation: $[\alpha]_D^{20} = -39.9^\circ$ (c=1, methanol).

J. (-)-cis-Terephthalic acid N,N-diisopropyl-N'-[3-(3-ethoxy-4-methoxyphenyl)-1-methylpiperidin-4-yl]diamide

CH: $C_{29}H_{41}N_3O_4$; MW: 495.67; yield: 91% of theory; m.p.: 75-82°C (unsharp range, solidified foam);

Optical rotation: $[\alpha]_D^{20} = -60.1^\circ$ (c=1, methanol).

K. (-)-cis-Terephthalic acid N,N-diisopropyl-N'-[3-(4-ethoxy-3-methoxyphenyl)-1-methylpiperidin-4-yl]diamide

EF: $C_{29}H_{41}N_3O_4$; MW: 495.67; yield: 89% of theory; m.p.: 72-80°C (unsharp range, solidified foam).

L. (-)-cis-Terephthalic acid N,N-dibutyl-N'-[3-(3-ethoxy-4-methoxyphenyl)-1-methylpiperidin-4-yl]diamide

EF: $C_{31}H_{45}N_3O_4$; MW: 523.73; yield: 90% of theory; m.p.: 98-102°C.

M. (-)-cis-4-(Hexahydroazepine-1-carbonyl)-N-[3-(3-ethoxy-4-methoxyphenyl)-1-piperidin-4-yl]benzamide

EF: C₂₉H₃₉N₃O₄; MW: 493.66; yield: 98% of theory; m.p.: 63-66°C (unsharp range, solidified foam).

The following compounds are obtained analogously to the process described in DE 42 17 401 when the corresponding 4-ethoxy-3-methoxy or 3-ethoxy-4-methoxy compounds are employed in the examples described there:

N. (-)-cis-4-Amino-3-(4-ethoxy-3-methoxyphenyl)-1-methylpiperidine dihydrochloride

EF: C₁₅H₂₄N₂O₂ × 2 HCl × 0.32 H₂O; MW: 343.06; m.p.: 241-243°C;

Optical rotation: $[\alpha]_D^{20} = -59.5^\circ$ (c=1, methanol).

O. (-)-cis-4-Amino-3-(3-ethoxy-4-methoxyphenyl)-1-methylpiperidine dihydrochloride

EF: C₁₅H₂₄N₂O₂ × 2 HCl × 0.96 H₂O; MW: 354.52; m.p.: 252-254°C;

Optical rotation: $[\alpha]_D^{20} = -65.5^\circ$ (c=1, methanol).

Commercial utility

The compounds according to the invention have valuable pharmacological properties which make them commercially utilizable. As selective inhibitors of type 3 and 4 of cyclic nucleotide phosphodiesterase (PDE3, PDE4), they are suitable on the one hand as bronchial therapeutics (for the treatment of airway obstructions on account of their dilating and cilium-stimulating action but also on account of their respiratory rate- and respiratory drive-increasing action), but on the other hand especially for the treatment of disorders of inflammatory nature, e.g. of the airways (asthma prophylaxis), of the skin, of the intestine, of the eyes and of the joints, which are mediated by mediators such as interferons, members of the tumor necrosis factor family, interleukins, chemokines, colony-stimulating factors, growth factors, lipid mediators (e.g., *inter alia*, PAF, platelet-activating factor), bacterial factors (e.g. LPS), immunoglobulins, oxygen free radicals and related free radicals (e.g. nitrogen monoxide NO), biogenic amines (e.g. histamine, serotonin), kinins (e.g. bradykinin), neurogenic mediators (such as substance P, neurokinin), proteins such as, for example, granular contents of leukocytes (*inter alia* cationic proteins of eosinophils) and adherent proteins (e.g. integrins). The compounds according to the invention have smooth muscle-relaxant action, e.g. in the region of the bronchial system, of the blood circulation, and of the efferent urinary passages. Furthermore they have a cilium-frequency increasing action, e.g. in the bronchial system.

In this context, the compounds according to the invention are distinguished by low toxicity, good human acceptance, good enteral absorption and high bioavailability, great therapeutic breadth, the absence of significant side effects and good water solubility.

On account of their PDE-inhibiting properties, the compounds according to the invention can be employed as therapeutics in human and veterinary medicine, where they can be used, for example, for the treatment and prophylaxis of the following diseases: acute and chronic (in particular inflammatory and allergen-induced) airway disorders of various origin (bronchitis, allergic bronchitis, bronchial asthma); disorders with a reduction of the cilium activity or with increased demands on the ciliar clearance (bronchitis, mucoviscidose); dermatoses (especially of proliferative, inflammatory and allergic type) such as, for example, psoriasis (vulgaris), toxic and allergic contact eczema, atopic eczema, seborrheic eczema, lichen simplex, sunburn, pruritis in the anogenital area, alopecia areata, hypertrophic scars, discoid lupus erythematosus, follicular and widespread pyodermias, endogenous and exogenous acne, acne rosacea and other proliferative, inflammatory and allergic skin disorders; disorders which are based on excessive release of TNF and leukotrienes, i.e., for example, disorders of the arthritis type (rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis and other arthritic conditions), systemic lupus erythematosus, disorders of the immune system (AIDS), including AIDS-related encephalopathies, autoimmune disorders such as diabetes mellitus (Type I, autoimmune diabetes), multiple sclerosis and of the type virus-, bacteria- or parasite-induced demyelination diseases, cerebral

malaria or Lyme's disease, shock symptoms [septic shock, endotoxin shock, Gram-negative sepsis, toxic shock syndrome and ARDS (adult respiratory distress syndrome)] and also generalized inflammations in the gastrointestinal region (Crohn's disease and ulcerative colitis); disorders which are based on allergic and/or chronic, faulty immunological reactions in the region of the upper airways (pharynx, nose) and of the adjacent regions (paranasal sinuses, eyes), such as, for example, allergic rhinitis/sinusitis, chronic rhinitis/sinusitis, allergic conjunctivitis and also nasal polyps; and also disorders of the central nervous system such as memory disorders and Alzheimer's disease, candidiasis, leishmaniases and leprosy.

On account of their vasorelaxant activity, the compounds according to the invention can also be used for the treatment of high blood pressure disorders of various origin such as, for example, pulmonary high blood pressure and the concomitant symptoms associated therewith, for the treatment of erectile dysfunction or colics of the kidneys and the ureters in connection with kidney stones.

On account of their cAMP-increasing action, however, they can also be used for disorders of the heart which can be treated by PDE inhibitors, such as, for example, cardiac insufficiency, and also as anti-thrombotic, platelet aggregation-inhibiting substances.

The invention further relates to a method for the treatment of mammals including humans who are suffering from one of the abovementioned diseases. The method comprises administering a therapeutically effective and pharmacologically tolerable amount of one or more of the compounds according to the invention to the sick mammal.

The invention further relates to the compounds according to the invention for use in the treatment and/or prophylaxis of the diseases mentioned.

The invention also relates to the use of the compounds according to the invention for the production of medicaments which are employed for the treatment and/or prophylaxis of the diseases mentioned.

The invention furthermore relates to medicaments for the treatment and/or prophylaxis of the diseases mentioned and which contain one or more of the compounds according to the invention.

Advantageously, the substances according to the invention are also suitable for combination with other substances which bring about stimulation of cAMP, such as prostaglandins (PGE₂, PGI₂ and prostacyclin) and their derivatives, direct adenylate cyclase stimulators such as forskolin and related substances, or substances indirectly stimulating adenylate cyclase, such as catecholamines and adrenergic receptor agonists, in particular beta mimetics. In combination, on account of their cAMP degradation-inhibiting action, they in this case display a synergistic, superadditive activity. This comes to bear,

for example, in their use in combination with PGE2 for the treatment of pulmonary hypertension.

The medicaments are prepared by methods known per se familiar to the person skilled in the art. As medicaments, the compounds according to the invention (= active compounds) are either employed as such, or preferably in combination with suitable pharmaceutical auxiliaries, e.g. in the form of tablets, coated tablets, capsules, suppositories, patches, emulsions, suspensions, gels or solutions, the active compound content advantageously being between 0.1 and 95%.

The person skilled in the art is familiar on the basis of his expert knowledge with the auxiliaries which are suitable for the desired pharmaceutical formulations. Beside solvents, gel-forming agents, ointments bases and other active compound excipients, it is possible to use, for example, antioxidants, dispersants, emulsifiers, preservatives, solubilizers or permeation promoters.

For the treatment of disorders of the respiratory tract, the compounds according to the invention are preferably also administered by inhalation. For this purpose, these are administered either directly as a powder (preferably in micronized form) or by atomization of solutions or suspensions which contain them. With respect to the preparations and administration forms, reference is made, for example, to the details in European Patent 163 965.

For the treatment of dermatoses, the compounds according to the invention are used in particular in the form of those medicaments which are suitable for topical application. For the production of the medicaments, the compounds according to the invention (= active compounds) are preferably mixed with suitable pharmaceutical auxiliaries and additionally processed to give suitable pharmaceutical formulations. Suitable pharmaceutical formulations which may be mentioned are, for example, powders, emulsions, suspensions, sprays, oils, ointments, fatty ointments, creams, pastes, gels or solutions.

The medicaments according to the invention are prepared by methods known per se. The dosage of the active compounds takes place in the order of magnitude customary for PDE-inhibitors. Thus topical application forms (such as, for example, ointments) for the treatment of dermatoses contain the active compounds in a concentration of, for example, 0.1-99%. The dose for administration by inhalation is customarily between 0.01 and 10 mg per spray burst. The customary dose in the case of systemic therapy (p.o. or i.v.) is between 0.1 and 200 mg per administration.

Biological investigations

In the investigation of PDE4 inhibition at the cellular level, the activation of inflammatory cells has particular importance. An example which may be mentioned is the FMLP (N-formylmethionylleucyl-phenylalanine)-induced superoxide production of neutrophilic granulocytes, which can be measured as luminol-potentiated chemoluminescence [McPhail LC, Strum SL, Leone PA and Sozzani S, The neutrophil respiratory burst mechanism. In "Immunology Series" 1992, 57, 47-76; ed. Coffey RG (Marcel Dekker, Inc., New York-Basel-Hong Kong)].

Substances which inhibit chemoluminescence and cytokine secretion and the secretion of inflammation-increasing mediators in inflammatory cells, in particular neutrophilic and eosinophilic granulocytes, are those which inhibit PDE4. This isoenzyme of the phosphodiesterase families is particularly represented in granulocytes. Its inhibition leads to an increase in the intracellular cyclic AMP concentration and thus to the inhibition of cellular activation. PDE4 inhibition by the substances according to the invention is thus a central indicator of the suppression of inflammatory processes. (Giembycz MA, Could isoenzyme-selective phosphodiesterase inhibitors render bronchodilatory therapy redundant in the treatment of bronchial asthma? Biochem Pharmacol 1992, 43, 2041-2051; Torphy TJ et al., Phosphodiesterase inhibitors: new opportunities for treatment of asthma. Thorax 1991, 46, 512-523; Schudt C et al., Zardaverine: a cyclic AMP PDE3/4 inhibitor. In "New Drugs for Asthma Therapy", 379-402, Birkhäuser Verlag Basel 1991; Schudt C et al., Influence of selective phosphodiesterase inhibitors on human neutrophil functions and levels of cAMP and Ca. Naunyn-Schmiedeberg's Arch Pharmacol 1991, 344, 682-690; Nielson CP et al., Effects of selective phosphodiesterase inhibitors on polymorphonuclear leukocytes respiratory burst. J Allergy Clin Immunol 1990, 86, 801-808; Schade et al., The specific type 3 and 4 phosphodiesterase inhibitor zardaverine suppress formation of tumor necrosis factor by macrophages. European Journal of Pharmacology 1993, 230, 9-14).

A. Methodology

1. Inhibition of PDE isoenzymes

The PDE activity was determined according to Thompson et al. (1) with some modifications (2). The test samples contained 40 mM tris HCl (pH 7.4), 5 mM $MgCl_2$, 0.5 μ M cAMP or cGMP, [3H] cAMP or [3H]cGMP (about 50,000 cpm/sample), the PDE isoenzyme-specific additions described in greater detail below, the indicated concentrations of inhibitor and an aliquot of the enzyme solution in a total sample volume of 200 μ l. Stock solutions of the compounds to be investigated in DMSO were prepared in concentrations such that the DMSO content in the test samples did not exceed 1% by volume - to avoid an effect on the PDE activity. After preincubation at 37°C for 5 minutes, the reaction was started by addition of the substrate (cAMP or cGMP). The samples were incubated at 37°C for a further 15 min. The reaction was terminated by addition of 50 μ l of 0.2N HCl. After cooling on ice for 10 minutes and addition of 25 μ g of 5'-nucleotidase (snake venom from *Crotalus atrox*), the mixture was again incubated at 37°C for 10 min and the samples were then applied to QAE Sephadex A-25 columns. The columns were eluted with 2 ml of 30 mM ammonium formate (pH 6.0). The radioactivity of the eluate was measured and corrected by the corresponding blank values. The proportion of hydrolyzed nucleotide in no case exceeded 20% of the original substrate concentration.

PDE1 (Ca^{2+} /calmodulin-dependent) from bovine brain: the inhibition of this isoenzyme was investigated in the presence of Ca^{2+} (1 mM) and calmodulin (100 nM) using cGMP as a substrate (3).

PDE2 (cGMP-stimulated) from rat hearts was purified chromatographically [Schudt et al. (4)] and investigated in the presence of cGMP (5 μ M) using cAMP as a substrate.

PDE3 (cGMP-inhibited) and PDE5 (cGMP-specific) were investigated in homogenates of human blood platelets [Schudt et al. (4)] using cAMP or cGMP as a substrate.

PDE4 (cAMP-specific) was investigated in the cytosol of human polymorphonuclear leukocytes (PMNL) [isolated from leukocyte concentrates, see Schudt et al. (5)] using cAMP as a substrate. The PDE3 inhibitor motapizone (1 μ M) was used in order to suppress the PDE3 activity emanating from contaminating blood platelets.

2. Inhibition of the formation of reactive oxygen species in human PMNL

The formation of reactive oxygen species determined with the aid of luminol-potentiated chemiluminescence (5) and the isolation of the PMNL from human blood (6) was carried out essentially as de-

- 27 -

scribed in (5) and (6): equal-size portions (0.5 ml) of the cell suspension (10^7 cells/ml) were preincubated at 37°C for 5 min in the absence or presence of the compounds to be investigated in a buffer solution comprising 140 mM NaCl, 5 mM KCl, 10 mM HEPES, 1 mM $\text{CaCl}_2/\text{MgCl}_2$, 1 mg/ml of glucose, 0.05% (w/v) BSA (bovine serum albumin), 10 μM luminol and 4 μM microperoxidase. Stock solutions of the compounds to be investigated in DMSO were prepared in concentrations such that the DMSO content in the test samples did not exceed 0.1% by volume - to avoid an effect on PDE activity. After preincubation, the test samples were transferred into the measuring apparatus ["Multi-Biolumat" LB 9505C from Berthold (Wildbad, Germany)] before stimulation with the receptor agonist FMLP (N-formylmethionylleucyl phenylalanine, 100 nM). The chemiluminescence was recorded continuously for 3 min; from this the AUC values were calculated.

3. Statistics

The IC_{50} values were determined from the concentration-inhibition curves by nonlinear regression using the GraphPad InPlot™ program (GraphPad Software Inc., Philadelphia, USA).

4. References

- (1) Thompson W.J. and Appleman M.M., Assay of cyclic nucleotide phosphodiesterase and resolution of multiple molecular forms of the enzyme; *Adv. Cycl. Nucl. Res.* **1979**, 10, 69-92
- (2) Bauer A.C. and Schwabe U., An improved assay of cyclic 3',5'-nucleotide phosphodiesterase with QAE Sephadex A-25; *Naunyn-Schmiedeberg's Arch. Pharmacol.* **1980**, 311, 193-198
- (3) Gietzen K., Sadorf I. and Bader H., A model for the regulation of the calmodulin-dependent enzymes erythrocyte Ca^{2+} -transport ATPase and brain phosphodiesterase by activators and inhibitors; *Biochem. J.* **1982**, 207, 541-548.
- (4) Schudt C., Winder S., Müller B. and Ukena D., Zardaverine as a selective inhibitor of phosphodiesterase isoenzymes; *Biochem. Pharmacol.* **1991**, 42, 153-162
- (5) Schudt C., Winder S., Forderkunn S., Hatzelmann A. and Ullrich V., Influence of selective phosphodiesterase inhibitors on human neutrophil functions and levels of cAMP and Ca; *Naunyn-Schmiedeberg's Arch. Pharmacol.* **1991**, 344, 682-690
- (6) Hatzelmann A. and Ullrich V., Regulation of 5-lipoxygenase activity by the glutathione status in human polymorphonuclear leukocytes; *Eur. J. Biochem.* **1987**, 169, 175-184

B. Results


In Table 1 below, the inhibitory concentrations determined according to Section A1 [inhibitory concentrations as $-\log IC_{50}$ (mol/l)] for the compounds according to the invention are indicated for various PDE isoenzymes. The numbers of the compounds correspond to the numbers of the examples.

Table 1

Compound	PDE5	PDE4	PDE3	PDE2	PDE1
	[-log IC ₅₀ , mol/l]				
3		6.45	7.14		
4	5.45	7.54	6.67	4.80	<4
5		7.75	7.15		
11		7.85	7.23		
16		7.96	6.73		
17		7.94	6.38		
18		7.87	6.74		
19		8.18	7.56		
21		7.67	6.34		
23		8.56	6.64		
24		8.51	7.64		

Patent Claims

1. Compounds of the formula I



in which

R1 is 1-4C-alkyl,

R2 is hydroxyl, 1-4C-alkoxy, 3-7C-cycloalkoxy, 3-7C-cycloalkylmethoxy or 1-4C-alkoxy which is completely or predominantly substituted by fluorine,

R3 is hydroxyl, 1-4C-alkoxy, 3-7C-cycloalkoxy, 3-7C-cycloalkylmethoxy or 1-4C-alkoxy which is completely or predominantly substituted by fluorine,

or in which

R2 and R3 together are a 1-2C-alkylenedioxy group,

R4 is a phenyl radical which is substituted by R5 and R6, where

R5 is hydrogen, hydroxyl, halogen, nitro, 1-4C-alkyl, trifluoromethyl or 1-4C-alkoxy,

R6 is CO-R7 or CO-R8, where

R7 is hydroxyl, 1-8C-alkoxy, 3-7C-cycloalkoxy or 3-7C-cycloalkylmethoxy and

R8 is N(R81)R82, where R81 and R82 independently of one another are hydrogen, 1-7C-alkyl, 3-7C-cycloalkyl or 3-7C-cycloalkylmethyl, or where R81 and R82, together and including the nitrogen atom to which both are bonded, are a 1-pyrrolidinyl, 1-piperidyl, 1-hexahydroazepinyl or 4-morpholinyl radical,

and the salts of these compounds.

2. Compounds of the formula I according to claim 1, in which

R1 is 1-4C-alkyl,

R2 is 1-4C-alkoxy, 3-7C-cycloalkoxy, 3-7C-cycloalkylmethoxy or 1-2C-alkoxy which is completely or predominantly substituted by fluorine,

R3 is 1-4C-alkoxy, 3-7C-cycloalkoxy, 3-7C-cycloalkylmethoxy or 1-2C-alkoxy which is completely or predominantly substituted by fluorine,

R4 is a phenyl radical which is substituted by R5 and R6, where

R5 is hydrogen, hydroxyl, halogen, nitro, 1-4C-alkyl, trifluoromethyl or 1-4C-alkoxy.

- R6 is CO-R7 or CO-R8, where
R7 is hydroxyl, 1-8C-alkoxy, 3-7C-cycloalkoxy or 3-7C-cycloalkylmethoxy and
R8 is N(R81)R82, where R81 and R82 independently of one another are hydrogen, 1-7C-alkyl, 3-7C-cycloalkyl or 3-7C-cycloalkylmethyl, or where R81 and R82, together and including the nitrogen atom to which both are bonded, are a 1-piperidyl, 1-hexahydroazepinyl or 4-morpholinyl radical,
and the salts of these compounds.

3. Compounds of the formula I according to claim 1, in which

- R1 is methyl,
R2 is 1-4C-alkoxy, 3-7C-cycloalkoxy or 1-2C-alkoxy which is completely or predominantly substituted by fluorine,
R3 is 1-4C-alkoxy, 3-7C-cycloalkoxy or 1-2C-alkoxy which is completely or predominantly substituted by fluorine,
R4 is a phenyl radical which is substituted by R5 and R6, where
R5 is hydrogen, hydroxyl, 1-4C-alkyl or 1-4C-alkoxy,
R6 is CO-R7 or CO-R8, where
R7 is hydroxyl, 1-8C-alkoxy or 3-7C-cycloalkoxy and
R8 is N(R81)R82, where R81 and R82 independently of one another are hydrogen, 1-7C-alkyl or 3-7C-cycloalkyl, or where R81 and R82, together and including the nitrogen atom to which both are bonded, are a 1-piperidyl, 1-hexahydroazepinyl or 4-morpholinyl radical,
and the salts of these compounds.

4. Compounds of the formula I according to claim 1, in which

- R1 is methyl,
R2 is methoxy or ethoxy,
R3 is methoxy or ethoxy,
R4 is a phenyl radical which is substituted by R5 and R6, where
R5 is hydrogen,
R6 is CO-R7 or CO-R8, where
R7 is hydroxyl or 1-8C-alkoxy and
R8 is N(R81)R82, where R81 and R82 independently of one another are hydrogen or 1-4C-alkyl or 5-7C-cycloalkyl, or where R81 and R82, together and including the nitrogen atom to which both are bonded, are a 1-piperidyl, 1-hexahydroazepinyl or 4-morpholinyl radical,
and the salts of these compounds.

5. Compounds of the formula I according to claim 1, in which

- R1 is methyl,

R2 is ethoxy,
 R3 is methoxy or ethoxy,
 R4 is a phenyl radical which is substituted by R5 and R6, where
 R5 is hydrogen,
 R6 is CO-R7 or CO-R8, where
 R7 is 1-4C-alkoxy and
 R8 is N(R81)R82, where R81 and R82 independently of one another are 1-4C-alkyl or 5-7C-cycloalkyl, or where R81 and R82, together and including the nitrogen atom to which both are bonded, are a 1-piperidyl or 1-hexahydroazepinyl radical,
 and the salts of these compounds.

6. Compounds of the formula I according to claim 1, in which the hydrogen atoms in positions 4a and 10b are in the cis position relative to one another, or the salts of these compounds.

7. Compounds of the formula I according to claim 1, which in the positions 4a and 10b have the same absolute configuration as the compound (-)-cis-4-amino-3-(3,4-dimethoxyphenyl)-1-methylpiperidine dihydrochloride having the optical rotation $[\alpha]_D^{22} = -57.1^\circ$ (c=1, methanol), which can be employed as a starting material.

8. A compound selected from the group consisting of

cis-8,9-Diethoxy-6-(4-isopropoxycarbonylphenyl)-2-methyl-1,2,3,4,4a,10b-hexahydrobenzo[c][1,6]-naphthyridine;

cis-8,9-Diethoxy-6-(4-diisopropylaminocarbonylphenyl)-2-methyl-1,2,3,4,4a,10b-hexahydrobenzo[c][1,6]naphthyridine;

cis-8,9-Diethoxy-6-(4-N-cyclohexyl-N-isopropylaminocarbonylphenyl)-2-methyl-1,2,3,4,4a,10b-hexahydrobenzo[c][1,6]naphthyridine;

cis-8,9-Diethoxy-6-(4-dibutylaminocarbonylphenyl)-2-methyl-1,2,3,4,4a,10b-hexahydrobenzo[c][1,6]naphthyridine;

cis-8,9-Diethoxy-6-[4-(hexahydroazepin-1-ylcarbonyl)phenyl]-2-methyl-1,2,3,4,4a,10b-hexahydrobenzo[c][1,6]naphthyridine;

cis-8,9-Diethoxy-6-[4-(piperidin-1-ylcarbonyl)phenyl]-2-methyl-1,2,3,4,4a,10b-hexahydrobenzo[c][1,6]naphthyridine;

cis-9-Ethoxy-8-methoxy-6-(4-diisopropylaminocarbonylphenyl)-2-methyl-1,2,3,4,4a,10b-hexahydrobenzo[c][1,6]naphthyridine;

cis-9-Ethoxy-8-methoxy-6-(4-dibutylaminocarbonylphenyl)-2-methyl-1,2,3,4,4a,10b-hexahydrobenzo[c][1,6]naphthyridine;

cis-9-Ethoxy-8-methoxy-6-[4-(hexahydroazepin-1-yl-carbonyl)phenyl]-2-methyl-1,2,3,4,4a,10b-hexa-

hydrobenzo[c][1,6]naphthyridine;

cis-9-Ethoxy-8-methoxy-6-[4-(piperidin-1-ylcarbonyl)phenyl]-2-methyl-1,2,3,4,4a,10b-hexahydrobenzo[c][1,6]naphthyridine

and the salts of these compounds, which can be prepared starting from (-)-cis-4-Amino-3-(3,4-diethoxyphenyl)-1-methylpiperidine dihydrochloride or (-)-cis-4-Amino-3-(3-ethoxy-4-methoxy)-1-methylpiperidine dihydrochloride.

9. A medicament comprising one or more compounds according to claim 1 together with the customary pharmaceutical auxiliaries and/or excipients.

10. The use of compounds according to claim 1 for the production of medicaments for the treatment of airway disorders and/or dermatoses.

INTERNATIONAL SEARCH REPORT

Inter. .nal Application No

PCT/EP 97/06096

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D471/04 A61K31/47 //(C07D471/04,221:00,221:00)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 91 17991 A (BYK GULDEN LOMBERG CHEMISCHE FABRIK) 28 November 1991 cited in the application see page 12-15; claim 1 ---	1-10
Y	EP 0 247 971 A (SANDOZ) 2 December 1987 cited in the application see page 1-10 ---	1-10
A	US 3 899 494 A (SANDOZ) 12 August 1975 cited in the application see column 6, line 63-68; claim 1 -----	1-10



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

9 February 1998

Date of mailing of the international search report

27. 02.98

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Lauro, P

INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern. Patent Application No

PCT/EP 97/06096

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9117991 A	28-11-91	AU 651592 B	28-07-94
		AU 7870191 A	10-12-91
		CA 2082923 A	17-11-91
		CN 1056691 A,B	04-12-91
		CS 9101447 A	18-03-92
		DE 4191109 A	01-04-93
		EP 0528922 A	03-03-93
		IL 98133 A	11-11-94
		US 5346904 A	13-09-94

EP 247971 A	02-12-87	AU 599981 B	02-08-90
		AU 7343787 A	03-12-87
		DK 273187 A	30-11-87
		JP 62294614 A	22-12-87

US 3899494 A	12-08-75	CH 571517 A	15-01-76
		CH 575413 A	14-05-76
		CH 571521 A	15-01-76
		CH 572055 A	30-01-76
		US 3966938 A	29-06-76
		AT 326674 B	29-12-75
		BE 767031 A	01-10-71
		CA 970377 A	01-07-75
		CH 527196 A	31-08-72
		CS 162753 B	15-07-75
		CS 167202 B	29-04-76
		DE 2123328 A	02-12-71
		DK 134488 B	15-11-76
		FR 2100649 A	24-03-72
		GB 1361441 A	24-07-74
		GB 1361442 A	24-07-74
		NL 7106502 A	16-11-71
		SE 387640 B	13-09-76
		CH 527822 A	15-09-72
		AU 6177573 A	24-04-75
		BE 806532 A	25-04-74
		DD 107457 A	05-08-74
		DE 2352909 A	09-05-74
		FR 2204418 A	24-05-74
		JP 49076900 A	24-07-74

Information on patent family members

PCT/EP 97/06096

Form PCT/ISA/210 (patent family annex) (July 1992)

THIS PAGE BLANK (USPTO)